



# Trenolin® Bouquet<sup>PLUS</sup>


Glycosidase for flavour release in must and wine

## Product description

Trenolin® Bouquet<sup>PLUS</sup> is a depsidase (cinnamyl esterase) free, aroma-releasing special enzyme.

The newly developed  $\beta$ -glycosidase activity can be used in mash, in must, during malolactic fermentation and in wine. Due to the sugar tolerant  $\beta$ -glycosidase activity, Trenolin® Bouquet<sup>PLUS</sup> can be applied before alcoholic fermentation to release aroma precursors.

Other benefits	<ul style="list-style-type: none"> <li>• Release of aromatic compounds from grapes</li> <li>• Release of various terpenes</li> <li>• Broad spectrum of uses, resulting in targeted and optimised increase of fruitiness</li> </ul>
Recommended for	<ul style="list-style-type: none"> <li>• All types of wine</li> </ul>
Advise	<ul style="list-style-type: none"> <li>• Enzyme purity is decisive particularly for aroma enzymes. Trenolin® Bouquet<sup>PLUS</sup> preserves released aroma compounds, as it is free of cinnamyl esterase activity.</li> </ul>



Permitted according to EU Commission Regulation no. 2019/934. User must check compliance with national regulations. Laboratory tested for purity and quality.

## PLUS - new formula means additional benefit!

The following properties are characteristic of the Trenolin®<sup>PLUS</sup> series of enzymes:

- Recipe optimised for modern wine-making
- Depsidase-free (free from cinnamyl esterase activity)
- No preservatives

## Dosage

	mL/100 L or 100 kg mash		
Mash	Approx. 7 - 10	During fermentation	Approx. 5 - 10
Must	Approx. 7 - 10	Wine	Approx. 10 - 15
Application time	Usually at least 1 week.		
Tip	Effectiveness depends on the quantity of product, temperature and reaction time. The treatment temperature should be more than 12 °C and preferably more than 15 °C. The higher the temperature, the more effective the enzyme. The natural upper limit is 55 °C.		
Application	Dilute the appropriate quantity of enzyme per tank with a little liquid to achieve better and more even distribution.		
Attention!	Bentonites deactivate the enzyme, so bentonite should be racked off before the addition of Trenolin® Bouquet <sup>PLUS</sup> or it should be used one week after the addition.		

## Storage

Store in a cool place. Packs which have been opened should be immediately tightly sealed and used up as soon as possible.