



# Trenolin® FastFlow

Enzyme for pressing, clarification and filtration

## Product description

Trenolin® FastFlow causes very rapid and intensive pectin degradation in mash and must, even with particularly pectin-rich grape varieties. The pressing yield and subsequent filtration performance of the wines are greatly improved.

Trenolin® FastFlow is capable of removing difficult to degrade pectin fractions (hairy regions) through the arabinogalactan-II-hydrolase (AG-II-hydrolase) activity. Trenolin® FastFlow is depectidase (cinnamyl esterase) free and therefore does not form undesirable volatile phenols.

Permitted according to EU Commission Regulation no. 2019/934. User must check compliance with national regulations. Laboratory tested for purity and quality.

## Treatment aim

Other benefits	<ul style="list-style-type: none"> <li>• Rapid, intensive pectin degradation in pectin-rich grape variety mashes such as Silvaner and Gewürztraminer. Marked improvement in press performance.</li> <li>• Pectin degradation from 8 °C on possible.</li> <li>• Targeted degradation of pectin side chains through AG-II-hydrolase for more effective pectin hydrolysis in juice, resulting faster loss of pectin's water-binding ability, rapid viscosity reduction and promotion of juice clarification.</li> <li>• Faster clarification when using LittoFresh® Origin, LittoFresh® ChitoFlot, e.IsingClair and KlarSol Super for sedimentation. Improved flotation processes by reducing neutral pectin fractions.</li> <li>• Increased filtration rates in the latter wines.</li> </ul>
Recommended for	<ul style="list-style-type: none"> <li>• Pectin-rich white and red grape varieties</li> </ul>

## Dosage

Application	Mash/Juice	Application time
Pectin-rich white grape varieties 8 - 15 °C	6 - 10 mL/100 kg or 100 L	2 - 4 hours
Red grape varieties up to 25 °C.	4 - 8 mL/100 kg or 100 L	during mash fermentation
Red grape varieties from 45 °C	3 - 6 mL/100 kg or 100 L	1 - 2 hours
Tip	Effective pectin degradation possible from 8 °C on. The natural upper limit is 55 °C.	
Application	Dilute the appropriate quantity of enzyme per tank with a little liquid to achieve a better and more even distribution.	
Attention!	Bentonites reduce enzyme content, so bentonite should only be used after the pectin has been fully degraded (pectin test).	

## Storage

Store in a cool place. Packs which have been opened should be tightly sealed and used up as soon as possible.

ERBSLÖH Geisenheim GmbH • Erbslöhstraße 1 • 65366 Geisenheim, Germany  
Tel.: +49 6722 708-0 • Fax: +49 6722 6098 • [info@erbsloeh.com](mailto:info@erbsloeh.com) • [www.erbsloeh.com](http://www.erbsloeh.com)

**ERBSLÖH**  
Fortschritt macht Zukunft®